

Package: utest (via r-universe)

May 24, 2026

Title Implementation of the Lind/Mehlum Utest

Version 0.4.0

Description An implementation of Lind and Mehlum's (2010) Utest to test for the presence of a U shaped or inverted U shaped relationship between variables in (generalized) linear models. It also implements a test of upward/downward sloping relationships at the lower and upper boundary of the data range.

License MIT + file LICENSE

Encoding UTF-8

LazyData true

Roxygen list(markdown = TRUE)

RoxygenNote 7.3.2

Suggests testthat

Repository <https://jotlind.r-universe.dev>

Date/Publication 2025-07-31 12:02:36 UTC

RemoteUrl <https://github.com/jotlind/utest>

RemoteRef HEAD

RemoteSha 28d2813cedec335428243230707bffce05af9250

Contents

uslopes	2
utest	3

Index	4
--------------	----------

uslopes

Estimates slopes at extremes

Description

This function estimates the slope of the relationship at the extreme points of the independent variable

Usage

```
uslopes(lmObject, vars, .vcov = NULL, x.min = NULL, x.max = NULL)
```

Arguments

lmObject	The model to be tested
vars	A vector with the name of the linear and squared terms. Can also be provided as a formula
.vcov	The covariance matrix to use
x.min	Lower bound of interval. If NULL, the minimum observed in the data is used.
x.max	Upper bound of interval. If NULL, the maximum observed in the data is used.

Details

The function computes slopes of a quadratic relationship at the lower and upper bound defined by `x.min` and `x.max`. Standard errors of the estimated slopes, t-values, and p-values from a one-sided test of a flat relationship as well as the extreme point of the estimated relationship are also provided.

Examples

```
x <- runif(100,min=-1,max=1)
xsq <- x^2
y <- x^2+rnorm(100)
mod <- lm(y~x+xsq)

uslopes(mod,c("x", "xsq"))
uslopes(mod,~x+xsq,x.max=0.8)
```

utest	<i>Perform Utest</i>
-------	----------------------

Description

This function computes the Lind/Mehlum test of a U shaped relationship.

Usage

```
utest(lmObject, vars, .vcov = NULL, x.min = NULL, x.max = NULL)
```

Arguments

lmObject	The model to be tested
vars	A vector with the name of the linear and squared terms. Can also be provided as a formula
.vcov	The covariance matrix to use
x.min	Lower bound of interval. If NULL, the minimum observed in the data is used.
x.max	Upper bound of interval. If NULL, the maximum observed in the data is used.

Details

To test for a U shaped or inverse U shaped relationship, it is necessary to provide an interval where the shape is located. A U shaped relationship is downward sloping at the lower bound and upward sloping at the upper bound, and vice versa for an inverted U shape.

The function assumes inputs from a model including a squared specification. The sign of the squared term is used to differentiate between U shaped and inverted U shaped relationships.

The function provides a test of the joint hypothesis of a downward sloping relationship at x.min and an upward sloping relationship at x.max for U shaped relationships and vice versa for inverted U shaped relationships, as detailed in Lind and Mehlum (2010).

References

J. T. Lind and H. Mehlum (2010), [With or without U? The appropriate test for a U shaped relationship](#). *Oxford Bulletin of Economics and Statistics* **72(1)**: 109-18.

Examples

```
x <- runif(100,min=-1,max=1)
xsq <- x^2
y <- x^2+rnorm(100)
mod <- lm(y~x+xsq)

utest(mod,c("x","xsq"))
utest(mod,~x+xsq,x.max=0.8)

mod.logit <- glm((y>0)~x+xsq,family="binomial")
utest(mod.logit,c("x","xsq"))
```

Index

uslopes, [2](#)
utest, [3](#)